

Discovery Research

In-Vitro ADME Capabilities



Discovery Research SUVEN Life Sciences Ltd

Serene Chambers, Road-5, Avenue-7, Banjara Hills,
Hyderabad-500034, India.

Contacts: @suven.com, nvsrk@suven.com

Overview



S. No	Title	Page No
1	Capabilities	3
2	Solubility and Distribution coefficient	4
3	Unbound fraction	5
4	P-glycoprotein (P-gp) Substrate assessment	6-7
5	Phenotyping	8
6	Enzyme Kinetics and CYP Inhibition	9
7	Time Dependent Inhibition	10
8	CYP Induction	11
9	Uptake Transporter	12
10	Quality Assurance	13
11	Contacts	14



Capabilities

Absorption	Solubility (Kinetic), Lipophilicity (LogD7.4), Permeability (PAMPA and Caco-2) and Efflux assays (Caco-2)
Distribution	Protein binding (plasma, brain homogenate, and microsomes), Blood to plasma partitioning (Species: rat, dog, and human)
Metabolism	Metabolic stability and intrinsic Clearance (S9, microsomes, rP450, hepatocytes), Reaction phenotyping (CYP and non-CYP, Enzyme kinetics, Chemical inhibition method and RAF), Metabolite identification and GSH trapping
Drug-Drug Interactions	Direct inhibitor (IC_{50} and K_i), Time dependent inhibition (Single point, IC_{50} shift, K_I and K_{inact} , Dialysis) CYP Induction (mRNA expression and enzyme activity) Uptake transporters: substrate and inhibition assays (OAT1, OAT3, OCT1, OCT2, OATP1B1, OATP1B3, MATE-1 and MATE-2K)



Solubility and Distribution coefficient

Kinetic solubility can help interpret the complication arising from the compound precipitation during biochemical, functional, and cell based assays
Identify poor soluble compounds that reduce productivity in drug discovery and development

Kinetic solubility

Phosphate buffer 7.4

Simulated gastric fluid

Simulated intestinal fluid


Lipophilicity is a key factor in determining the permeation of physiological membrane, protein binding, and target affinity

Log D 7.4

Miniaturized shake-flask method

n-Octanol / Phosphate buffer 7.4


Cyclohexane / Phosphate buffer 7.4






Unbound fraction (plasma/brain homogenate/microsomes)

Knowledge of Unbound fraction is essential to correct the total plasma concentration and correlate with efficacy



Brain homogenate




Rapid Equilibrium Dialysis (RED) and High Throughput (HT) Dialysis



P-glycoprotein (P-gp) Substrate assessment

A net flux ratio (or efflux ratio (ER)) of ≥ 2 in cells that express P-gp (e.g., Caco-2 cells) suggest that New chemical entity is a P-gp substrate


Test System:	Caco-2
Test Conc.:	10 μ M
Incubation:	60 min, 37 °C
Positive control	Digoxin
Membrane	TEER & Lucifer Yellow
Analysis	LC-MS/MS





P-glycoprotein (P-gp) Inhibitor assessment

Test System	Caco-2
Test Concentration	50 μ M
Positive control	Verapamil
Negative Control	Phenelzine
P-gp Substrate	Digoxin
Incubation	1 hour
Membrane Integrity	TEER & Lucifer Yellow
Analysis	LC-MS/MS






Phenotyping (Chemical inhibition method and RAF)

Contribution $\geq 25\%$ by an enzyme is considered significant based on in vitro phenotyping studies and Human Pharmacokinetic study

Evaluate the role of CYP1A2, CYP2B6, 121 CYP2C8, CYP2C9, CYP2C19, CYP2D6, and CYP3A4/5 and additional enzymes including MAO, FMO, and UGT

In vitro Phenotyping studies include chemical inhibition and Metabolism in recombinant enzymes (RAF approach)

P450	HLM	rCYP	CL-RAF ($\text{pmol}\cdot\text{mg}^{-1}$)	Mean CYP abundance ($\text{pmol}\cdot\text{mg}^{-1}$)	CL-ISEF
	CL_{int} ($\text{uL}\cdot\text{min}^{-1}\cdot\text{mg}^{-1}$)	CL_{int} ($\text{uL}\cdot\text{min}^{-1}\cdot\text{pmol CYP}^{-1}$)			
1A2	7 ± 0.3	1.2 ± 0.05^a	5.6 ± 0.5	39	0.14 ± 0.01
2A6	1298 ± 58	23 ± 0.5	56.9 ± 3	27	2.11 ± 0.1
2B6	3 ± 0.2	0.13 ± 0.01	19.8 ± 1	16	1.24 ± 0.07
2C8	777 ± 38	13 ± 2	58.8 ± 6	22.4	2.62 ± 0.3
2C9	96 ± 6	4.3 ± 0.1	21.2 ± 1	61	0.35 ± 0.01
2C19	0.4 ± 0.1	0.15 ± 0.01	2.8 ± 0.4	11	0.25 ± 0.04
2D6	33 ± 0.3	32 ± 1	1.0 ± 0.03	12.6	0.08 ± 0.002
2 E1	8 ± 0.3	0.13 ± 0.001	62.0 ± 2	64.5	0.96 ± 0.03
3A4	399 ± 33	18 ± 1.3	22 ± 1	93	0.24 ± 0.01





Enzyme Kinetics and CYP Inhibition

CYP Inhibition (IC_{50} and K_i)
 CYP1A2, CYP2B6, 121 CYP2C8, CYP2C9, CYP2C19, CYP2D6, and CYP3A4/5

Basic Model (I/ K_i) or Mechanistic Static Model to predict drug interaction risk


Validated marker activities for major CYP isoforms in HLM and rCYP


P450	HLM			rCYP		
	K_m (μM)	V_{max} ($pmol \cdot min^{-1} \cdot mg^{-1}$)	Kinetic Mechanism	K_m μM	V_{max} ($pmol \cdot min^{-1} \cdot pmolCYP^{-1}$)	Kinetic Mechanism
1A2	59 ± 3	395 ± 10	MM	82 ± 5.9^a	31 ± 2.7	NC
2A6	0.5 ± 0.07	649 ± 10	MM	0.5 ± 0.004	11.4 ± 0.3	MM
2B6	70 ± 3	187 ± 7	SI	88.4 ± 10.6	11.9 ± 0.5	SI
2C8	2.7 ± 0.2	2097 ± 51	MM	0.5 ± 0.1	6.6 ± 0.1	MM
2C9	9.7 ± 0.5	933 ± 17	MM	1.1 ± 0.04	5.0 ± 0.1	MM
2C19	92 ± 8	38 ± 2	MM	13.4 ± 0.7	2.0 ± 0.01	MM
2D6	4.6 ± 0.1	152 ± 4	MM	0.2 ± 0.1	6.3 ± 0.1	MM
2E1	177 ± 9	1384 ± 61	MM	101.5 ± 2.2	12.8 ± 0.1	MM
3A4	3.0 ± 0.2	1197 ± 63	MM	3.1 ± 0.2	56.1 ± 0.9	SI

^a- S_{50} instead of K_m ; ^b- Negative co-operativity, clearance read from plot of $v/[S]$ vs $[S]$;




Time Dependent Inhibition





Fold change relative to control
(Mean \pm SEM)




CYP Induction evaluated by “Gold standard method” using cryopreserved plateable hepatocytes from three donors at three concentrations of new chemical entity with both mRNA expression and enzyme activities monitored using RT-PCR and LC-MS/MS respectively. Vehicle control, Positive control, and Negative controls are included in the assay



Uptake Transporter (IC₅₀ assay)

Transporter	Substrate	Positive control	Absolute IC ₅₀ (uM)
OAT1	Para-amino hippuric acid (PAH)	Flufenamic acid	0.5
OAT3	Estrone 3-sulfate (E3S)	Indomethacin	0.7
OCT1	Tetraethyl ammonium (TEA)	Verapamil	5.0
OCT2	Metformin	Imipramine	6.4
OATP1B1	Estradiol β-D Glucuronide	Sulfasalazine	3.4
OATP1B3	Estradiol β-D Glucuronide	Rifampicin	2.3



Compound ID	OCT2 Abs IC ₅₀ (µM)	Literature Reported
Imipramine	6.4	3.3
Tacrine	3.4	3.1



- Independent Quality Assurance team
- Quality System Procedures (QSP's) for Quality System Management and Standard Operating Procedures (SOP's) for Operation, Calibration, Maintenance of Equipment's
- Document and Data Control, Conducting Internal Audits, Study Specific Audits
- Dedicated Archive facility for the retention of the records
- Facility audited and approved by many global pharmaceutical companies and majority of Indian Pharma Companies

Contacts



Venkat Jasti

Chairman & CEO

E-mail: info@suven.com

Ramakrishna Nirogi

Vice President, Discovery Research

E-mail: nvsrk@suven.com